

METHOD OF DIAGNOSING,
MONITORING, STAGING, IMAGING AND TREATING PROSTATE CANCER

FIELD OF THE INVENTION

This invention relates, in part, to newly developed
5 assays for detecting, diagnosing, monitoring, staging,
prognosticating, imaging and treating cancers, particularly
prostate cancer.

BACKGROUND OF THE INVENTION

Cancer of the prostate is the most prevalent malignancy
10 in adult males, excluding skin cancer, and is an increasingly
prevalent health problem in the United States. In 1996, it
was estimated that 41,400 deaths would result from this
disease in the United States alone, indicating that prostate
cancer is second only to lung cancer as the most common cause
15 of death in the same population. If diagnosed and treated
early, when the cancer is still confined to the prostate, the
chances of cure is significantly higher.

Treatment decisions for an individual are linked to the
stage of prostate cancer present in that individual. A common
20 classification of the spread of prostate cancer was developed
by the American Urological Association (AUA). The AUA system
divides prostate tumors into four stages, A to D. Stage A,
microscopic cancer within prostate, is further subdivided into
sub-stages A1 and A2. Sub-stage A1 is a well-differentiated
25 cancer confined to one site within the prostate. Treatment
is generally observation, radical prostatectomy, or radiation.
Sub-stage A2 is a moderately to poorly differentiated cancer
at multiple sites within the prostate. Treatment is radical
prostatectomy or radiation. Stage B, palpable lump within the

09807200-052904
F005250-00240860

1. The first step is to identify the problem or question that needs to be addressed. This involves understanding the context and the specific requirements of the task.

5 Treatment for sub-stages B1 and B2 is either radical prostatectomy or radiation. Stage C is a large cancer mass involving most or all of the prostate and is also further subdivided into two stages. In sub-stage C1, the cancer forms a continuous mass that may have extended beyond the prostate.

10 In sub-stage C2, the cancer forms a continuous mass that invades the surrounding tissue. Treatment for both these sub-stages is radiation with or without drugs to address the cancer. The fourth stage, Stage D is metastatic cancer and is also subdivided into two sub-stages. In sub-stage D1, the

15 cancer appears in the lymph nodes of the pelvis. In sub-stage D2, the cancer involves tissues beyond lymph nodes. Treatment for both of these sub-stages is systemic drugs to address the cancer as well as pain.

Accordingly, there is a great need for increasingly sensitive methods for diagnosing and staging of prostate cancer in a human patient to determine whether or not such cancer has metastasized and for monitoring the progress of a cancer which has not metastasized for the onset of metastasis.

- 3 -

In the present invention, methods are provided for detecting, diagnosing, monitoring, staging, prognosticating, imaging and treating prostate cancer via the cancer specific gene referred to herein as CSG. CSG refers, among other things, to native protein expressed by the gene comprising the polynucleotide sequence of SEQ ID NO:1. The amino acid sequence of a polypeptide encoded by SEQ ID NO:1 is depicted herein as SEQ ID NO:2. In the alternative, what is meant by CSG as used herein, means the native mRNA encoded by the gene comprising the polynucleotide sequence of SEQ ID NO:1 or levels of the gene comprising the polynucleotide sequence of SEQ ID NO:1.

Other objects, features, advantages and aspects of the present invention will become apparent to those of skill in the art from the following description. It should be understood, however, that the following description and the specific examples, while indicating preferred embodiments of the invention are given by way of illustration only. Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following description and from reading the other parts of the present disclosure.

SUMMARY OF THE INVENTION

Toward these ends, and others, it is an object of the present invention to provide a method for diagnosing the presence of prostate cancer by analyzing for changes in levels of CSG in cells, tissues or bodily fluids compared with levels of CSG in preferably the same cells, tissues, or bodily fluid type of a normal human control, wherein a change in levels of

- 4 -

CSG in the patient versus the normal human control is associated with prostate cancer.

Further provided is a method of diagnosing metastatic prostate cancer in a patient having prostate cancer which is not known to have metastasized by identifying a human patient suspected of having prostate cancer that has metastasized; analyzing a sample of cells, tissues, or bodily fluid from such patient for CSG; comparing the CSG levels in such cells, tissues, or bodily fluid with levels of CSG in preferably the same cells, tissues, or bodily fluid type of a normal human control, wherein an increase in CSG levels in the patient versus the normal human control is associated with prostate cancer which has metastasized.

Also provided by the invention is a method of staging prostate cancer in a human which has such cancer by identifying a human patient having such cancer; analyzing a sample of cells, tissues, or bodily fluid from such patient for CSG; comparing CSG levels in such cells, tissues, or bodily fluid with levels of CSG in preferably the same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in CSG levels in the patient versus the normal human control is associated with a cancer which is progressing and a decrease in the levels of CSG is associated with a cancer which is regressing or in remission.

Further provided is a method of monitoring prostate cancer in a human having such cancer for the onset of metastasis. The method comprises identifying a human patient having such cancer that is not known to have metastasized; periodically analyzing a sample of cells, tissues, or bodily fluid from such patient for CSG; comparing the CSG levels in such cells, tissue, or bodily fluid with levels of CSG in

- 5 -

preferably the same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in CSG levels in the patient versus the normal human control is associated with a cancer which has metastasized.

5 Further provided is a method of monitoring the change in stage of prostate cancer in a human having such cancer by looking at levels of CSG in a human having such cancer. The method comprises identifying a human patient having such cancer; periodically analyzing a sample of cells, tissues, or
10 bodily fluid from such patient for CSG; comparing the CSG levels in such cells, tissue, or bodily fluid with levels of CSG in preferably the same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in CSG levels in the patient versus the normal human control is
15 associated with a cancer which is progressing and a decrease in the levels of CSG is associated with a cancer which is regressing or in remission.

Further provided are antibodies targeted against CSG or fragments of such antibodies which can be used to detect or
20 image localization of CSG in a patient for the purpose of detecting or diagnosing a disease or condition. Such antibodies can be polyclonal, monoclonal, or omniclonal or prepared by molecular biology techniques. The term "antibody", as used herein and throughout the instant
25 specification is also meant to include aptamers and single-stranded oligonucleotides such as those derived from an in vitro evolution protocol referred to as SELEX and well known to those skilled in the art. Antibodies can be labeled with a variety of detectable labels including, but not limited to,
30 radioisotopes and paramagnetic metals. These antibodies or fragments thereof can also be used as therapeutic agents in

T063530"0020350

2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590 2591 2592 2593 2594 2595 2596 2597 2598 2599 2600 2601 2602 2603 2604 2605 2606 2607 2608 2609 2610 2611 2612 2613 2614 2615 2616 2617 2618 2619 2620 2621 2622 2623 2624 2625 2626 2627 2628 2629 2630 2631 2632 2633 2634 2635 2636 2637 2638 2639 2640 2641 2642 2643 2644 2645 2646 2647 2648 2649 2650 2651 2652 2653 2654 2655 2656 2657 2658 2659 2660 2661 2662 2663 2664 2665 2666 2667 2668 2669 2670 2671 2672 2673 2674 2675 2676 2677 2678 2679 2680 2681 2682 2683 2684 2685 2686 2687 2688 2689 2690 2691 2692 2693 2694 2695 2696 2697 2698 2699 2700 2701 2702 2703 2704 2705 2706 2707 2708 2709 2710 2711 2712 2713 2714 2715 2716 2717 2718 2719 2720 2721 2722 2723 2724 2725 2726 2727 2728 2729 2730 2731 2732 2733 2734 2735 2736 2737 2738 2739 2740 2741 2742 2743 2744 2745 2746 2747 2748 2749 2750 2751 2752 2753 2754 2755 2756 2757 2758 2759 2760 2761 2762 2763 2764 2765 2766 2767 2768 2769 2770 2771 2772 2773 2774 2775 2776 2777 2778 2779 2780 2781 2782 2783 2784 2785 2786 2787 2788 2789 2790 2791 2792 2793 2794 2795 2796 2797 2798 2799 2800 2801 2802 2803 2804 2805 2806 2807 2808 2809 2810 2811 2812 2813 2814 2815 2816 2817 2818 2819 2820 2821 2822 2823 2824 2825 2826 2827 2

15 DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to diagnostic assays and methods, both quantitative and qualitative for detecting, diagnosing, monitoring, staging and prognosticating cancers by comparing levels of CSG in a human patient with those of CSG in a normal human control. What is meant by levels of CSG as used herein, means levels of the native protein expressed by the gene comprising the polynucleotide sequence of SEQ ID NO:1. The amino acid sequence of a polypeptide encoded by SEQ ID NO:1 is depicted herein as SEQ ID NO:2. The native protein being detected, may be whole, a breakdown product, a complex of molecules or chemically modified. In the alternative, what is meant by levels of CSG as used herein, means levels of the native mRNA encoded by the gene comprising the polynucleotide sequence of SEQ ID NO:1 or levels of DNA comprising the polynucleotide sequence of SEQ ID NO:1. Such levels are

- 7 -

preferably measured in at least one of cells, tissues and/or bodily fluids, including determination of normal and abnormal levels. Thus, for instance, a diagnostic assay in accordance with the invention for diagnosing over-expression of CSG protein compared to normal control bodily fluids, cells, or tissue samples may be used to diagnose the presence of prostate cancer.

All the methods of the present invention may optionally include measuring the levels of other cancer markers as well as CSG. Other cancer markers, in addition to CSG, useful in the present invention will depend on the cancer being tested and are known to those of skill in the art.

Diagnostic Assays

The present invention provides methods for diagnosing the presence of prostate cancer by analyzing for changes in levels of CSG in cells, tissues or bodily fluids compared with levels of CSG in cells, tissues or bodily fluids of preferably the same type from a normal human control, wherein an increase in levels of CSG in the patient versus the normal human control is associated with the presence of prostate cancer.

Without limiting the instant invention, typically, for a quantitative diagnostic assay a positive result indicating the patient being tested has cancer is one in which cells, tissues or bodily fluid levels of the cancer marker, such as CSG, are at least two times higher, and most preferably are at least five times higher, than in preferably the same cells, tissues or bodily fluid of a normal human control.

The present invention also provides a method of diagnosing metastatic prostate cancer in a patient having prostate cancer which has not yet metastasized for the onset of metastasis. In the method of the present invention, a

Without limiting the instant invention, typically, for a quantitative diagnostic assay a positive result indicating the cancer in the patient being tested or monitored has metastasized is one in which cells, tissues or bodily fluid levels of the cancer marker, such as CSG, are at least two times higher, and most preferably are at least five times higher, than in preferably the same cells, tissues or bodily fluid of a normal patient.

- 9 -

Normal human control as used herein includes a human patient without cancer and/or non cancerous samples from the patient; in the methods for diagnosing or monitoring for metastasis, normal human control may preferably also include
5 samples from a human patient that is determined by reliable methods to have prostate cancer which has not metastasized.

Staging

The invention also provides a method of staging prostate cancer in a human patient. The method comprises identifying
10 a human patient having such cancer and analyzing a sample of cells, tissues or bodily fluid from such human patient for CSG. The measured CSG levels in the patient are then compared with levels of CSG in preferably the same cells, tissues or bodily fluid type of a normal human control, wherein an
15 increase in CSG levels in the human patient versus the normal human control is associated with a cancer which is progressing and a decrease in the levels of CSG (but still increased over true normal levels) is associated with a cancer which is regressing or in remission.

20 Monitoring

Further provided is a method of monitoring prostate cancer in a human patient having such cancer for the onset of metastasis. The method comprises identifying a human patient having prostate cancer that is not known to have metastasized;
25 periodically analyzing cells, tissues or bodily fluid from such human patient for CSG; comparing the CSG levels in such cells, tissues or bodily fluid with levels of CSG in preferably the same cells, tissues or bodily fluid type of a normal human control, wherein an increase in CSG levels in the
30 human patient versus the normal human control is associated

- 10 -

with a cancer which has metastasized. In this method, normal human control samples may also include prior patient samples.

Further provided by this invention is a method of monitoring the change in stage of prostate cancer in a human patient having such cancer. The method comprises identifying a human patient having prostate cancer; periodically analyzing cells, tissues or bodily fluid from such human patient for CSG; comparing the CSG levels in such cells, tissues or bodily fluid with levels of CSG in preferably the same cells, tissues or bodily fluid type of a normal human control, wherein an increase in CSG levels in the human patient versus the normal human control is associated with a cancer which is progressing in stage and a decrease in the levels of CSG is associated with a cancer which is regressing in stage or in remission. In this method, normal human control samples may also include prior patient samples.

Monitoring patients for onset of metastasis is periodic and preferably done on a quarterly basis. However, monitoring may be performed more or less frequently depending on the cancer, the particular patient, and the stage of the cancer.

Assay Techniques

Assay techniques that can be used to determine levels of gene expression (including protein levels), such as CSG of the present invention, in a sample derived from a patient are well known to those of skill in the art. Such assay methods include, without limitation, radioimmunoassays, reverse transcriptase PCR (RT-PCR) assays, immunohistochemistry assays, *in situ* hybridization assays, competitive-binding assays, Western Blot analyses, ELISA assays and proteomic approaches: two-dimensional gel electrophoresis (2D electrophoresis) and non-gel based approaches such as mass

- 11 -

spectrometry or protein interaction profiling. Among these, ELISAs are frequently preferred to diagnose a gene's expressed protein in biological fluids.

An ELISA assay initially comprises preparing an antibody, 5 if not readily available from a commercial source, specific to CSG, preferably a monoclonal antibody. In addition a reporter antibody generally is prepared which binds specifically to CSG. The reporter antibody is attached to a detectable reagent such as radioactive, fluorescent or 10 enzymatic reagent. For example, detectable agents such as horseradish peroxidase enzyme and alkaline phosphatase are routinely used in these types of assays.

To carry out the ELISA, antibody specific to CSG is incubated on a solid support, e.g. a polystyrene dish, that 15 binds the antibody. Any free protein binding sites on the dish are then covered by incubating with a non-specific protein such as bovine serum albumin. Next, the sample to be analyzed is incubated in the dish, during which time CSG binds to the specific antibody attached to the polystyrene 20 dish. Unbound sample is washed out with buffer. A reporter antibody specifically directed to CSG and linked to a detectable reagent such as horseradish peroxidase is placed in the dish resulting in binding of the reporter antibody to any monoclonal antibody bound to CSG. Unattached reporter 25 antibody is then washed out. Reagents for detection of peroxidase activity, including a colorimetric substrate are then added to the dish. Immobilized peroxidase linked to CSG antibodies produces a colored reaction product. The amount of color developed in a given time period is proportional to 30 the amount of CSG protein present in the sample. Quantitative

- 12 -

results typically are obtained by reference to a standard curve.

A competition assay can also be employed wherein antibodies specific to CSG are attached to a solid support. 5 Labeled CSG and a sample derived from the host are then passed over the solid support. The amount of label detected which is attached to the solid support can be correlated to a quantity of CSG in the sample.

Nucleic acid methods can be used to detect CSG mRNA as 10 a marker for prostate cancer. Polymerase chain reaction (PCR) and other nucleic acid methods, such as ligase chain reaction (LCR) and nucleic acid sequence based amplification (NASABA), can be used to detect malignant cells for diagnosis and monitoring of various malignancies. For example, reverse- 15 transcriptase PCR (RT-PCR) is a powerful technique which can be used to detect the presence of a specific mRNA population in a complex mixture of thousands of other mRNA species. In RT-PCR, an mRNA species is first reverse transcribed to complementary DNA (cDNA) with use of the enzyme reverse 20 transcriptase; the cDNA is then amplified as in a standard PCR reaction. RT-PCR can thus reveal by amplification the presence of a single species of mRNA. Accordingly, if the mRNA is highly specific for the cell that produces it, RT-PCR can be used to identify the presence of a specific type of 25 cell.

Hybridization to clones or oligonucleotides arrayed on a solid support (i.e. gridding) can be used to detect the expression of and quantitate the level of expression of that gene. In this approach, a cDNA encoding the CSG gene is fixed 30 to a substrate. The substrate may be of any suitable type including but not limited to glass, nitrocellulose, nylon or

- 13 -

plastic. At least a portion of the DNA encoding the CSG gene is attached to the substrate and then incubated with the analyte, which may be RNA or a complementary DNA (cDNA) copy of the RNA isolated from the tissue of interest.

5 Hybridization between the substrate bound DNA and the analyte can be detected and quantitated by several means including, but not limited to, radioactive labeling or fluorescence labeling of the analyte or a secondary molecule designed to detect the hybrid. Quantitation of the level of gene
10 expression can be done by comparison of the intensity of the signal from the analyte compared with that determined from known standards. The standards can be obtained by *in vitro* transcription of the target gene, quantitating the yield, and then using that material to generate a standard curve.

15 Of the proteomic approaches, 2D electrophoresis is a technique well known to those in the art. Isolation of individual proteins from a sample such as serum is accomplished using sequential separation of proteins by different characteristics usually on polyacrylamide gels.
20 First, proteins are separated by size using an electric current. The current acts uniformly on all proteins so that smaller proteins move farther on the gel than larger proteins. The second dimension applies a current perpendicular to the first and separates proteins not on the basis of size but on
25 the specific electric charge carried by each protein. Since no two proteins with different sequences are identical on the basis of both size and charge, the result of a 2D separation is a square gel in which each protein occupies a unique spot. Analysis of the spots with chemical or antibody probes or
30 subsequent protein microsequencing can reveal the relative

09807200-052904
FOUO

- 14 -

abundance of a given protein and the identity of the proteins in the sample.

The above tests can be carried out on samples derived from a variety of cells, bodily fluids and/or tissue extracts including both homogenates or solubilized tissue obtained from a human patient. Tissue extracts can be obtained from tissue biopsy and autopsy material. Bodily fluids useful in the present invention include blood, urine, saliva or any other bodily secretion or derivative thereof. By blood it is meant to include whole blood, plasma, serum or any derivative of blood.

In Vivo Antibody Use

Antibodies which specifically bind to CSG can also be used in vivo in patients suspected of suffering from prostate cancer. Specifically, antibodies which specifically bind a CSG can be injected into a patient suspected of having prostate cancer for diagnostic and/or therapeutic purposes. The preparation and use of antibodies for in vivo diagnosis is well known in the art. For example, antibody-chelators labeled with Indium-111 have been described for use in the radioimmunoscentigraphic imaging of carcinoembryonic antigen expressing tumors (Sumerdon et al. Nucl. Med. Biol. 1990 17:247-254). In particular, these antibody-chelators have been used in detecting tumors in patients suspected of having recurrent colorectal cancer (Griffin et al. J. Clin. Onc. 1991 9:631-640). Antibodies with paramagnetic ions as labels for use in magnetic resonance imaging have also been described (Lauffer, R.B. Magnetic Resonance in Medicine 1991 22:339-342). Antibodies directed against CSG can be used in a similar manner. Labeled antibodies which specifically bind CSG can be injected into patients suspected of having prostate

- 15 -

cancer for the purpose of diagnosing or staging of the disease status of the patient. The label used will be selected in accordance with the imaging modality to be used. For example, radioactive labels such as Indium-111, Technetium-99m or Iodine-131 can be used for planar scans or single photon emission computed tomography (SPECT). Positron emitting labels such as Fluorine-19 can be used in positron emission tomography. Paramagnetic ions such as Gadolinium (III) or Manganese (II) can be used in magnetic resonance imaging (MRI). Localization of the label permits determination of the spread of the cancer. The amount of label within an organ or tissue also allows determination of the presence or absence of cancer in that organ or tissue.

For patients diagnosed with prostate cancer, injection of an antibody which specifically binds CSG can also have a therapeutic benefit. The antibody may exert its therapeutic effect alone. Alternatively, the antibody can be conjugated to a cytotoxic agent such as a drug, toxin or radionuclide to enhance its therapeutic effect. Drug monoclonal antibodies have been described in the art for example by Garnett and Baldwin, Cancer Research 1986 46:2407-2412. The use of toxins conjugated to monoclonal antibodies for the therapy of various cancers has also been described by Pastan et al. Cell 1986 47:641-648. Yttrium-90 labeled monoclonal antibodies have been described for maximization of dose delivered to the tumor while limiting toxicity to normal tissues (Goodwin and Meares Cancer Supplement 1997 80:2675-2680). Other cytotoxic radionuclides including, but not limited to Copper-67, Iodine-131 and Rhenium-186 can also be used for labeling of antibodies against CSG.

- 16 -

Antibodies which can be used in these *in vivo* methods include polyclonal, monoclonal and omniclonal antibodies and antibodies prepared via molecular biology techniques. Antibody fragments and aptamers and single-stranded
5 oligonucleotides such as those derived from an *in vitro* evolution protocol referred to as SELEX and well known to those skilled in the art can also be used.

The present invention is further described by the following examples. These examples are provided solely to
10 illustrate the invention by reference to specific embodiments. These exemplifications, while illustrating certain aspects of the invention, do not portray the limitations or circumscribe the scope of the disclosed invention.

EXAMPLES

15 The examples are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. Routine molecular biology techniques of the following examples can be carried out as described in standard laboratory manuals, such as
20 Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989).

Example 1

Suppression subtractive hybridization (CLONTECH PCR-SELECT)

25 CLONTECH PCR-SELECT is a PCR based subtractive hybridization method designed to selectively enrich for cDNAs corresponding to mRNAs differentially expressed between two mRNA populations (Diatchenko et al, *Proc. Natl. Acad. Sci. USA*, Vol. 93, pp. 6025-6030, 1996). In this method,
30 differentially expressed mRNAs are enriched based on a selective amplification. cDNA is prepared from the two mRNA

090072000-050004

- 17 -

populations which are to be compared. These include the Tester population which is a cDNA population in which the differentially expressed messages are sought and the Driver population which is a cDNA population in which the differentially expressed transcripts are absent or low. The tester sample is separated in two parts and different PCR adapters are ligated to the 5' ends. Each tester is separately annealed to excess driver in the first annealing and then pooled and again annealed in the second annealing to excess driver. During the first annealing, sequences common to both populations anneal. Additionally the concentration of high and low abundance messages are normalized since annealing is faster for abundant molecules due to the second order kinetics of hybridization. During the second annealing cDNAs unique or overabundant to the tester can anneal together. Such molecules have different adapters at their ends. The addition of additional driver during the second annealing enhances the enrichment of the desired differentially expressed sequences. During subsequent PCR, molecules that have different adapters at each end amplify exponentially. Molecules which have identical adapters, or adapters at only one end, or no adapters (driver sequences) either do not amplify or undergo linear amplification. The end result is enrichment for cDNAs corresponding to differentially expressed messages unique to the tester or up regulated in the tester.

This technique was used to identify transcripts unique to cancer tissues or messages overexpressed in the cancer process. To do this, pairs of samples isolated from a cancer
30 tissue were used as the "tester", and non-cancer tissue as the "driver". The non-cancer "driver" can be from the same

- 18 -

individual and tissue as the tumor (Matched). Alternatively, the "driver" can be from a different individual but the same tissue as the tumor sample (unmatched). In some cases, the "driver" comprises mixtures of cDNAs derived from non-cancer tissues different from the cancer tissue. This approach allows the identification of transcripts whose expression is specific or up-regulated in the cancer tissue. Such transcripts may or may not be cancer specific in their expression.

10 Subtractive hybridization was carried out using as "tester" a mixture of three RNAs from human adenocarcinomas and as "driver" a mixture of RNAs from five human normal tissues (spleen, pancreas, heart, kidney and small intestine).

The subtracted mixture was cloned and two hundred clones were sequenced. One of the sequences matched Incyte clone ID 3966820. The electronic Northern for this clone showed the highest number of ESTs came from prostate, compared with other tissues (prostate 107 followed by uterus with 20).

The PCR-select clone detected a transcript of 1.9 kb by hybridization in Northern blots. Amongst 17 tissues tested prostate showed the highest abundance for this transcript.

Example 2: Relative Quantitation of Gene Expression

Real-Time quantitative PCR with fluorescent Taqman probes is a quantitation detection system utilizing the 5'-3' nuclease activity of Taq DNA polymerase. The method uses an internal fluorescent oligonucleotide probe (Taqman) labeled with a 5' reporter dye and a downstream, 3' quencher dye. During PCR, the 5'-3' nuclease activity of Taq DNA polymerase releases the reporter, whose fluorescence can then be detected

09807200-052904

- 19 -

by the laser detector of the Model 7700 Sequence Detection System (PE Applied Biosystems, Foster City, CA, USA).

Amplification of an endogenous control is used to standardize the amount of sample RNA added to the reaction and
5 normalize for Reverse Transcriptase (RT) efficiency. Either cyclophilin, glyceraldehyde-3-phosphate dehydrogenase (GAPDH) or 18S ribosomal RNA (rRNA) is used as this endogenous control. To calculate relative quantitation between all the samples studied, the target RNA levels for one sample were
10 used as the basis for comparative results (calibrator). Quantitation relative to the "calibrator" can be obtained using the standard curve method or the comparative method (User Bulletin #2: ABI PRISM 7700 Sequence Detection System).

The tissue distribution and the level of the target gene
15 were evaluated for every sample in normal and cancer tissues. Total RNA was extracted from normal tissues, cancer tissues, and from cancers and the corresponding matched adjacent tissues. Subsequently, first strand cDNA was prepared with reverse transcriptase and the polymerase chain reaction was
20 done using primers and Taqman probes specific to each target gene. The results were analyzed using the ABI PRISM 7700 Sequence Detector. The absolute numbers are relative levels of expression of the target gene in a particular tissue compared to the calibrator tissue.

25 Real-Time quantitative PCR was done using the following primers:

pro108 Reverse GCCTTCAGCCGTGGGTAGT (SEQ ID NO:3)

pro108 Forward GACAGCGGCTTCACCTTCTC (SEQ ID NO:4)

The absolute numbers depicted in Table 1 are relative
30 levels of expression of the CSG referred to herein as (pro108) (SEQ ID NO:1) in 11 normal different tissues. All the values

- 20 -

are compared to normal pancreas (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

5 Table 1: Relative Levels of CSG Pro108 Expression in Pooled Samples

10

15

Tissue	NORMAL
Colon Ascending	4.56
Kidney	0.78
Liver	5.66
Ovary	41.07
Small Intestine	0.78
Spleen	3.29
Stomach	11.39
Testis	2.04
Uterus	15.56
Pancreas	1.00
Prostate	2.32

The relative levels of expression in Table 1 show the
 20 highest mRNA expression in ovary (41.07) and uterus (15.56), two female specific tissues. Except for stomach (11.39) that shows high levels of mRNA for pro108, the other tissues including prostate show comparable low levels of expression.

The absolute numbers in Table 1 were obtained analyzing
 25 pools of samples of a particular tissue from different individuals. They can not be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 2.

30 The absolute numbers depicted in Table 2 are relative levels of expression of the CSG pro108 in 70 pairs of matching samples. All the values are compared to normal pancreas (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual.

T006250 00220860

- 21 -

Unmatched samples were used for prostatitis (prostatitis 1 and 2) and Benign Prostate Hyperplasia (BPH 1 to 6).

Table 2: Relative Levels of CSG Pro108 Expression in Individual Samples

5	Sample ID	Tissue	Cancer	Benign Diseases	Normal Adjacent Tissue
	Pro12B	Prostate 1	253.40		7.70
	Pro78XB	Prostate 2	1020.46		96.67
	Pro84XB	Prostate 3	2055.11		70.52
	Pro101XB	Prostate 4	872.61		218.58
10	Pro91X	Prostate 5	539.32		195.36
	Pro13XB	Prostate 6	0.60		0.49
	Pro23B	Prostate 7	929.30		747.00
	Pro90XB	Prostate 8	160.50		21.00
	Pro18XB	Prostate 9	45.63		31.49
15	Pro20XB	Prostate 10	312.50		20.00
	Pro34B	Prostate 11	1351.18		142.02
	Pro65XB	Prostate 12	1305.15		8.37
	Pro69XB	Prostate 13	486.50		14.50
	Pro10R	Prostatitis 1		122.45	
20	Pro20R	Prostatitis 2		83.38	
	Pro258BPH	BPH 1		39.33	
	Pro263CBPH	BPH 2		451.98	
	Pro267ABPH	BPH 3		90.78	
	Pro271ABPH	BPH 4		5.21	
25	Pro460ZBPH	BPH 5		15.64	
	ProC032BPH	BPH 6		46.85	
	Bld32XK	Bladder 1	2.86		3.88
	Bld46XK	Bladder 2	15.62		1.45

5	Bld66X	Bladder 3	32.67		19.90
	BldTR14	Bladder 4	330.60		235.50
	ClnAS67	Colon 1	71.26		0.51
	ClnRC01	Colon 2	5.08		25.90
	ClnSG45	Colon 3	19.09		6.61
	ClnSG67	Colon 4	65.12		20.61
	ClnTX01	Colon 5	116.97		6.25
	ClnB34	Colon 6	3.98		2.45
	ClnB56	Colon 7	1.98		1.41
	ClnC9XR	Colon 8	6.88		2.00
10	ClnCXGA	Colon 9	2.25		2.62
	ClnAS89	Colon 10	77.44		2.39
	ClnTX67	Colon 11	167.73		11.51
	End10479	Endometrium 1	7.63		26.67
	End8911	Endometrium 2	16.19		64.98
	End8963	Endometrium 3	12.14		65.89
	End5XA	Endometrium 4	99.39		23.51
	End65RA	Endometrium 5	6.94		16.06
	Kid109XD	Kidney 1	65.57		41.93
	Kid10XD	Kidney 2	39.67		9.71
20	Kid11XD	Kidney 3	12.68		1.62
	Kid126XD	Kidney 4	43.71		7.59
	Kid12XD	Kidney 5	2.50		70.80
	Kid5XD	Kidney 6	2.11		74.29
	Kid6XD	Kidney 7	19.36		0.72
	Liv42X	Liver 1	39.00		6.40
	Liv15XA	Liver 2	9.23		4.59
	Liv94XA	Liver 3	7.07		8.09

- 23 -

	LngAC66	Lung 1	47.30		16.20
	LngBR94	Lung 2	106.50		3.40
	LngLC109	Lung 3	9.60		34.10
	LngLC71	Lung 4	70.30		64.00
5	LngSQ56	Lung 5	12.40		53.80
	LngSQ79	Lung 6	108.40		77.20
	Lng60XL	Lung 7	0.55		1.47
	Lng75XC	Lung 8	2.34		3.11
	LngC17X	Lung 9	52.96		31.71
10	LngAC69	Lung 10	654.84		42.67
	LngC20X	Lung 11	2.82		0.22
	Mam47XP	Mammary Gland 1	3.40		1.89
	Mam82XI	Mammary Gland 2	33.29		4.44
	MamB011X	Mammary Gland 3	10.61		1.50
15	MamA06X	Mammary Gland 4	14.62		1.55
	Ovr	Ovary	10.00		8.00
	Pan77X	Pancreas 1	91.27		24.50
	Pan71XL	Pancreas 2	1.55		1.30
	Pan82XP	Pancreas 3	5.49		7.53
20	Pan92X	Pancreas 4	1069.00		688.40
	StoAC99	Stomach 1	50.70		18.60
	StoTA73	Stomach 2	37.70		130.30
	SmI21XA	Small Intestine 1	12.42		5.31
	SmIH89	Small Intestine 2	23.26		2.63
25	Tst39X	Testis	27.95		7.01

T06250"00220360

- 24 -

Utr85XU	Uterus 1	3.30		4.91
Utr23XU	Uterus 2	44.48		29.04

Table 2 shows the results for the analysis of 148 samples distributed in 14 different tissue types. Tables 1 and 2 represent a combined total of 159 samples in 16 different tissue types.

In the analysis of matching samples, the higher levels of expression were in prostate, showing a high degree of tissue specificity for prostate tissue. Of all the samples different from prostate which were analyzed (57 matching samples), only a few samples (lung 10 and pancreas 4) showed an expression comparable or higher than the median for the mRNA expression in prostate cancer (median: 539.32).

In addition, the level of mRNA expression in cancer samples was compared with the isogenic normal adjacent tissue from the same individual. This comparison provides an indication of specificity for the cancer stage (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 2 shows overexpression of the CSG prol08 in 13 of 13 prostate cancer tissues compared with their respective normal adjacent (prostate samples #1 to #13). Thus, there was overexpression in the cancer tissue for 100% of the prostate matching samples tested.

Altogether, the level of tissue specificity, plus the mRNA overexpression in 100% of the prostate matching samples tested are indicative of the CSG prol08 being a diagnostic marker for prostate cancer.